

Radiative B decays at LHCb

Fatima Soomro, Imperial College London


IOP 2010, 29-31 March 2010, University College London

Imperial College
London



- ✓ The LHCb detector (introduced by previous speakers)
- Introduction to radiative B decays
- Photon polarization in radiative B decays
- Photon polarization measurement at LHCb
 - systematics
- Conclusion and outlook

- Radiative B decays are $b \rightarrow s(d) \gamma$ transitions
- In the SM, allowed through a penguin loop
- Are sensitive to NP contribution
- Useful in constraining NP
 - ❑ Test mass scale of NP particles
 - ❑ Test the couplings of NP particles
 - ❑ Test if the couplings have a V-A structure




Branching ratio measurements

- $B_{\text{inclusive, exp}}(B \rightarrow X_s \gamma) = 3.56 \pm 0.26 \times 10^{-4}$ [1] [3]
- $B_{\text{inclusive, th}}(B \rightarrow X_s \gamma) = 3.15 \pm 0.23 \times 10^{-4}$ [2]

Good agreement puts limits on NP models

Exclusion areas in parameter space of a CMSSM model. Green: constraints from $b \rightarrow s \gamma$, brown: constraints from LSP, light blue: area favoured by WMAP [13]




In this talk, I will concentrate on measurements with $B \rightarrow X_s \gamma$, where X_s is a CP eigenstate (f^{CP}), in particular, $B_s \rightarrow \phi \gamma$.

Test the structure of NP operators if they contribute by measuring the photon polarization

In the SM, quarks that couple to the W are left handed


\Rightarrow the photons are predominantly L handed in \overline{B}^0 decays



Test the structure of NP operators if they contribute by measuring the photon polarization

In the SM, quarks that couple to the W are left handed

\Rightarrow the photons are predominantly L handed in \overline{B}^0 decays



In the SM the ratio of “wrong” helicity photons is $\propto m_s/m_b$ and predicted to be $\sim 0.4\%$

\Rightarrow in some NP scenarios can be up to 10% [4]

\Rightarrow for example: LR symmetric model and the unconstrained MSSM model, predict that it can be very large, without affecting the BR

(gluon emission can give upto a 1% effect as well [14])

In a $B \rightarrow f^{CP} \gamma$ decay, the time dependent decay width is parametrized as

$$\Gamma_{B_s \rightarrow f^{CP} \gamma}(t) = |A|^2 e^{-\Gamma_s t} \left(\cosh \frac{\Delta\Gamma_s t}{2} - \mathcal{A}^\Delta \sinh \frac{\Delta\Gamma_s t}{2} + \mathcal{C} \cos \Delta m_s t - \mathcal{S} \sin \Delta m_s t \right)$$

$$\Gamma_{\bar{B}_s \rightarrow f^{CP} \gamma}(t) = |A|^2 e^{-\Gamma_s t} \left(\cosh \frac{\Delta\Gamma_s t}{2} - \mathcal{A}^\Delta \sinh \frac{\Delta\Gamma_s t}{2} - \mathcal{C} \cos \Delta m_s t + \mathcal{S} \sin \Delta m_s t \right)$$

Where in the SM,

$$\mathcal{C} \approx 0, \quad \mathcal{S} \approx \sin 2\psi \sin \varphi_{(s)} \approx 0 \quad (\text{as } \varphi_{(s)} \text{ is small}),$$

$$\text{and } \mathcal{A}^\Delta \approx \sin 2\psi \cos \varphi_{(s)}$$

The parameter ψ contains information about the photon polarization:

$$\tan \psi \equiv \frac{\mathcal{A}(\bar{B}_s \rightarrow f^{CP} \gamma_R)}{\mathcal{A}(\bar{B}_s \rightarrow f^{CP} \gamma_L)}$$

Adding the two equations for the \bar{B}_s and B_s

$$\Gamma_{B_s \rightarrow f^{CP} \gamma}(t) = |A|^2 e^{-\Gamma_s t} \left(\cosh \frac{\Delta\Gamma_s t}{2} - \mathcal{A}^\Delta \sinh \frac{\Delta\Gamma_s t}{2} + \mathcal{C} \cos \Delta m_s t - \mathcal{S} \sin \Delta m_s t \right)$$

$$\Gamma_{\bar{B}_s \rightarrow f^{CP} \gamma}(t) = |A|^2 e^{-\Gamma_s t} \left(\cosh \frac{\Delta\Gamma_s t}{2} - \mathcal{A}^\Delta \sinh \frac{\Delta\Gamma_s t}{2} - \mathcal{C} \cos \Delta m_s t + \mathcal{S} \sin \Delta m_s t \right)$$


one gets $\Gamma_{B_s \rightarrow f^{CP} \gamma}(t) = |A|^2 e^{-\Gamma_s t} \left(\cosh \frac{\Delta\Gamma_s t}{2} - \mathcal{A}^\Delta \sinh \frac{\Delta\Gamma_s t}{2} \right)$


Adding the two equations for the \bar{B}_s and B_s

$$\Gamma_{B_s \rightarrow f^{CP} \gamma}(t) = |A|^2 e^{-\Gamma_s t} \left(\cosh \frac{\Delta\Gamma_s t}{2} - \mathcal{A}^\Delta \sinh \frac{\Delta\Gamma_s t}{2} + \mathcal{C} \cos \Delta m_s t - \mathcal{S} \sin \Delta m_s t \right)$$



$$\Gamma_{\bar{B}_s \rightarrow f^{CP} \gamma}(t) = |A|^2 e^{-\Gamma_s t} \left(\cosh \frac{\Delta\Gamma_s t}{2} - \mathcal{A}^\Delta \sinh \frac{\Delta\Gamma_s t}{2} - \mathcal{C} \cos \Delta m_s t + \mathcal{S} \sin \Delta m_s t \right)$$

one gets $\Gamma_{B_s \rightarrow f^{CP} \gamma}(t) = |A|^2 e^{-\Gamma_s t} \left(\cosh \frac{\Delta\Gamma_s t}{2} - \mathcal{A}^\Delta \sinh \frac{\Delta\Gamma_s t}{2} \right)$






Divide $A^\Delta = 0.0$ by $A^\Delta = 0.2$ distribution



- Can be done with untagged B_s sample,
 $\sigma_{A^\Delta} = 0.22$ (nominal LHCb year, 11 k signal events expected)
- For a tagged analysis, $\sigma_S = 0.1$ (nominal LHCb year) (remember $S \approx \sin 2\psi \sin \phi_s$)
 - measurements at Belle and BaBar with $B_d \rightarrow K^*(K_s \pi^0)\gamma$ $S = -0.19 \pm 0.23$ [12]
 - Belle and Babar analyses based on a few hundred signal events
- Experimental requirements (it is a tough measurement):
 - Proper time resolution/bias to be precisely known
 - The lifetime acceptance function to be precisely known
 - Background distribution in lifetime to be known

- Proper time resolution is a measure of how well the B_s lifetime is reconstructed



Fitted with a double gaussian

$$\mu_{\text{core}} = 3.9 \pm 0.6 \text{ fs (notice the unit)}$$

$$\sigma_{\text{core}} = 61 \pm 1 \text{ fs (fraction}_{\text{core}} = 85\%)$$

$$\mu_{\text{wide}} = 10.6 \pm 2 \text{ fs}$$

$$\sigma_{\text{wide}} = 154 \pm 3 \text{ fs}$$

- A bias of 5 fs gives an uncertainty on A^Δ which is 1/3 of the statistical uncertainty for 2fb^{-1} (nominal LHCb year)

The B_s proper time is given by

$$\tau = \frac{\vec{P} \cdot \vec{d}}{|\vec{P}|^2} m$$

Where \vec{P} is the momentum of the B_s ,
 \vec{d} is the distance between the primary and
the B_s decay vertex, and m is the B_s mass

The B_s proper time is given by


$$\tau = \frac{\vec{P} \cdot \vec{d}}{|\vec{P}|^2} m$$

Where \vec{P} is the momentum of the B_s ,

\vec{d} is the distance between the primary and the B_s decay vertex, and m is the B_s mass

Is dominated by the vertex reconstruction
the ϕ vertex becomes increasingly difficult to reconstruct as the \vec{P}_ϕ increases

Control channel: $B_s \rightarrow J/\psi \phi$



The B_s proper time is given by

$$\tau = \frac{\vec{P} \cdot \vec{d}}{|\vec{P}|^2} m$$



Where \vec{P} is the momentum of the B_s ,
 \vec{d} is the distance between the primary and
 the B_s decay vertex, and m is the B_s mass

Is dominated by the vertex reconstruction
 the ϕ vertex becomes increasingly difficult
 to reconstruct as the \vec{P}_ϕ increases


Control channel: $B_s \rightarrow J/\psi \phi$

Are dominated by the photon momentum
 reconstruction (ECAL resolution)

Control channel: $B_d \rightarrow K^* \gamma$



- The efficiency to reconstruct and select events as a function of their proper time



- To keep it as simple as possible, we try to avoid lifetime biasing cuts at the trigger and offline selection level
- Nevertheless, is non trivial
 - Calibrate using $B_s \rightarrow J/\psi \phi$

1. For the proper time bias:

- ❑ Look at the difference between the ϕ vertex and the J/ψ vertex in $B_s \rightarrow J/\psi\phi$
- ❑ Establish a γ momentum correction from $B_d \rightarrow K^*\gamma$ (by constraining the B_d mass)


2. For the proper time acceptance:

Again using $B_s \rightarrow J/\psi\phi$, compare the acceptance when cuts are applied to the J/ψ vertex and to the ϕ vertex

3. For the background model:


Study mass sidebands

- Radiative decays of B hadrons are sensitive probes of NP
 - A lot of measurements done by B factories using $B_d \rightarrow K^* \gamma$
 - The measurement of photon polarization (using $B_s \rightarrow \phi \gamma$) will require a lot of data and a very good control of systematics
- We already have a ϕ
 - Just need some highly energetic photons



- Radiative decays of B hadrons are sensitive probes of NP
 - A lot of measurements done by B factories using $B_d \rightarrow K^* \gamma$
 - The measurement of photon polarization (using $B_s \rightarrow \phi \gamma$) will require a lot of data and a very good control of systematics

- We already have a ϕ
 - Just need some highly energetic photons



- LHCb can make competitive measurements with early data (which I did not have the time to talk about)
 - World's best measurement of the direct CP asymmetry in $B_d \rightarrow K^* \gamma$
 - Ratio of BR $B_d \rightarrow K^* \gamma / B_s \rightarrow \phi \gamma$

References and backup slides

- [1] C.Amsler et al. [Particle Data Group], Phys. Lett. B 667, 1 (2008)
- [2] M.Misiak et al., Phys. Rev. Lett. 98, 022002 (2007)
- [3] The Belle Collaboration, Measurement of Inclusive Radiative B-meson Decays with a Photon Energy Threshold of 1.7 GeV, Phys. Rev. Lett.103:241801 (2009)
- [4] D. Atwood, M. Gronau and A. Soni, Phys. Rev. Lett. 79, 185 (1997)
- [5] The BaBar collaboration, Measurement of $B \rightarrow K^*(892)\gamma$ Branching Fractions and CP and Isospin Asymmetries, Phys. Rev. Lett. 103, 211802 (2009)
- [6] M. Matsumori, A. I. Sanda, and Y.-Y. Keum, Phys. Rev. D 72, 014013 (2005)
- [7] The LHCb Collaboration, B. Adeva et al, Roadmap for selected key measurements of LHCb, LHCb-PUB-2009-029, arXiv:0912.4179v1 [hep-ex]
- [8] Gautam Bhattacharyya, Gustavo C. Branco and Debajyoti Choudhury, Phys.Lett. B336 (1994) 487-493; Erratum-ibid. B340 (1994)
- [9] The BaBar collaboration, Phys. Rev. Lett. 98, 151802 (2007)
- [10] Michael Gronau, P and CP violation in B physics, Pramana62:255-268,2004
- [11] Jacques Lefrancois, Marie-H´elene Schune, LHCb-PUB-2009-008
- [13] John Ellis, Keith A. Olive, Yudi Santoso and Vassilis C. Spanos, Phys.Lett.B573:162-172,2003
- [14] arXiv:hep-ph/0609037

Experiment	$N(B\bar{B})$	$S_{CP}(b \rightarrow s\gamma)$	$C_{CP}(b \rightarrow s\gamma)$	Correlation
$K^*(892)\gamma$				
<i>BABAR</i> [340]	431M	$-0.08 \pm 0.31 \pm 0.05$	$-0.15 \pm 0.17 \pm 0.03$	0.05
Belle [341]	535M	$-0.32^{+0.36}_{-0.33} \pm 0.05$	$0.20 \pm 0.24 \pm 0.05$	0.08
Average		-0.19 ± 0.23	-0.03 ± 0.14	0.06
Confidence level		0.43 (0.8 σ)		
$K_S^0\pi^0\gamma$ (including $K^*(892)\gamma$)				
<i>BABAR</i> [342]	232M	-0.06 ± 0.37	-0.48 ± 0.22	0.05
Belle [341]	535M	$-0.10 \pm 0.31 \pm 0.07$	$0.20 \pm 0.20 \pm 0.06$	0.08
Average		-0.09 ± 0.24	-0.12 ± 0.15	0.06
Confidence level		0.08 (1.8 σ)		

[12] Heavy Flavor Averaging Group, arXiv:0808.1297v3 [hep-ex] (2009)

Branching Fraction and Photon Energy Spectrum for $b \rightarrow s\gamma$, *The CLEO collaboration*, Phys. Rev. Lett. 87, 251807 (2001)






FIG. 2. Observed laboratory frame photon energy spectrum (weights per 100 MeV) for on minus scaled off minus B backgrounds, the putative $b \rightarrow s\gamma$ plus $b \rightarrow d\gamma$ signal. No corrections have been applied for resolution or efficiency. Also shown is the spectrum from Monte Carlo simulation of the Ali-Greub spectator model with parameters $\langle m_b \rangle = 4.690$ GeV, $P_F = 410$ MeV/ c , a good fit to the data.



Sensitivity of the $B \rightarrow X_s \gamma$ branching ratio to the top-quark mass

Bound on the charged-Higgs mass in the type-II 2HDM derived from the analysis of the inclusive $B \rightarrow X_s \gamma$ branching ratio (solid line)



- $S_{K^*\gamma}$ is the time dependent CP asymmetry of $B \rightarrow K^*\gamma$. In 97 it was noticed that the standard model (V-A interaction !) produces mainly left handed photons in $\bar{B} \rightarrow \bar{K}^*\gamma_{L(R)}$, $O(m_s/m_b)$. A non-vanishing time dependent CP-asymmetry demands both chiralities and is therefore very small $\sim -2\%$. It was noticed that gluon emission from c and u-loops produce both chiralities and it was guesstimated to give a contribution of -10% ([hep-ph/0412019](http://arxiv.org/abs/hep-ph/0412019)) plus a large uncertainty. We have performed a straightforward calculation and have shown that the contribution is $0.5\% \pm 1\%$! ([hep-ph/0609037](http://arxiv.org/abs/hep-ph/0609037)) and therefore $S_{K^*\gamma}$ etc remain important (quasi) null tests of the standard model. The current HFAG value is $S_{K^*\gamma} = -28 \pm 26\%$! All TDCP asymmetries for $B \rightarrow V\gamma$ are observables of primary interest for the SuperB-factory
[<http://www.ippp.dur.ac.uk/~zwicky/>]

1. Direct CP and isospin asymmetry in $B \rightarrow K^* \gamma$
 - Results from experiment [5] agree with theory [6], but large errors
 - $-0.033 < A_{\text{cp}}(B_d \rightarrow K^* \gamma) < 0.028$ (theory: $< 1\%$)
 - $0.017 < \Delta_{0^-} < 0.116$ (theory: $\sim 4\%$)
 - LHCb can make a competitive measurement of $A_{\text{cp}}(B_d \rightarrow K^* \gamma)$ with only 100 pb^{-1} [7]
2. The photon energy spectrum in inclusive $b \rightarrow s \gamma$ decays is a useful experimental test of
 - The parton model, mass of spectator quark and the motion of the b quark inside the hadron [3]
3. Inclusive $b \rightarrow s \gamma$ and $b \rightarrow d \gamma$ [8,9]
 - Provide measurements of V_{ts} and V_{td} and their ratio
 - Compare to the ones measured in B_s and B_d oscillations
4. Photon polarization

1. $B_d \rightarrow K^* e e$
 - Angle b/w the $K\pi$ plane and ee plane
 - LHCb sensitivity to the the fraction of wrongly polarized photons is 0.1 for 2 fb^{-1} [[11](#)]
2. $\Lambda_b \rightarrow \Lambda (p\pi)\gamma$
 - A_{FB} of the proton flight direction wrt the Λ_b in Λ rest frame is proportional to the photon polarization
3. B to $h_1 h_2 h_3$ gamma (K resonances)
 - Only K(1400) has sensitivity, need to separate it from the other resonances
4. $B_s \rightarrow f^{cp}\gamma$