

Experimental realisation of the weak measurement process

- How do you do a strong and weak measurement.
- Two experiments using photons
 - Modified Stern-Gerlach
 - Youngs's 2-slit experiment
- Preliminary thoughts on using massive particles
 - Electrons
 - Neutrons
 - An other?
 - This is a work in progress

Experiments using weak measurement

The weak and strong measurements are carried out on non-commuting variables: Spin in the z and x axes; momentum and position.

There is a debate as to whether this is really a measurement process. I will illustrate that by looking at an experiment using a modified Stern-Gerlach where the usual spin measurement is amplified.

Then I will look at how the particle trajectories in a Young's 2-slit experiment can be mapped out.


Original Stern-Gerlach experiment

Silver atoms were originally used. They have one valence electron around a filled core and it behaved like a spin-1/2 particle.

The classical prediction was the beam should spread out in a continuous manner.

The observation is a beam split into two parts: spin-up, spin-down.

A STRONG MEASUREMENT



Reduce the magnetic field the beams are not well separated but overlapping.

A WEAK MEASUREMENT

Aharonov et al asked what would happen if you follow a weak measurement by a strong one (post-selection)?

Particle physicists use weak measurement

G. J. Feldman et al., Phys. Rev. Lett. 48, 66 (1982)

The weak measurement is a sum of overlapping Gaussians, each one centred on one of the values of the variable being measured.

If the entire distribution of events is mapped out then its centroid is the mean value of the variable being measured.

In the measurement of the τ -lifetime the position resolution of the detectors used was not good enough to resolve the decay length of the τ -lepton in a single event.

The “apparent decay length” has a very broad distribution, dominated by the position-measurement errors. With a large sample of events the life-time can be deduced from the mean (centroid) of the distribution.


Using a Double Stern-Gerlach apparatus

I. M. Duck and P. M. Stevenson, Phys. Rev. D, 1989

The particles enter along the y-axis with spin oriented at an angle α in the zx-plane.

The first S-G is oriented along the z-axis. **Weak**

The second S-G is oriented along the x-axis. **Strong**







$A_w = \lambda \tan(\alpha/2)$, $\lambda \propto$ magnetic moment of the particle.
Note what happens as α approaches π , A_w gets very large.

What happens as α approaches π ?

Let $\varepsilon = \pi - \alpha$.

Scale is $\sigma_x = 1$.



- (a) $\phi(p)$ resembles a Gaussian.
 (b) Close up shows the peak is at 5.


- (a) $\phi(p)$ is now antisymmetric.
 (b) The resulting probability distribution peaks at ± 70 .

Optical analogue of the Stern-Gerlach apparatus


Polarised light from a laser is used instead of spin-1/2 particles.

Polariser P and analyser A select the initial and final polarisations at angles α and β respectively.

A birefringent crystal provides the weak measuring device. It introduces a small lateral displacement between the o-ray and the e-ray.



Realisation of the optical analogue




Results of the optical analogue

The wavelength of the light $a=0.64\mu\text{m}$

(a) $\alpha=\beta=\pi/4$, corresponding to aligned polarizers. The measured intensity profile is the result of the constructive addition of two approximately Gaussian distributions.

(b) $\alpha=\pi/4$, $\beta=3\pi/4+0.022$, corresponding to a measurement of the weak value. The centroid of the distribution is shifted by $A_w=12\mu\text{m}=20a$.

(c) $\alpha=\pi/4$, $\beta=3\pi/4$, corresponding to crossed polarizers, or orthogonal initial and final states. The separation of the two peaks is $120a$.




Mapping trajectories of photons in 2-slit experiment

Kocsis et al, Science 332:1170, 2011

A quantum dot was used as a source of single photons. This made sure that only one photon was in the apparatus at a time.


The strong measurement of position is carried out by observing the fringe pattern at a range of horizontal positions




A 50:50 beam splitter.

Birefringent calcite is used to for the weak measurement of momentum. It imparts a small k_x -dependent phase shift ($p=hk$). Linear polarisation becomes slightly elliptical

Reconstruction of the trajectories



The magenta line is after constant background has been subtracted



**Preliminary thoughts on using
massive particles
(work in progress)**


Proposal

Carry out similar experiments using massive particles such as electrons and neutrons with a view of demonstrating weak measurement and exploring the existence and nature of the quantum potential .


The first step is to build a Double Stern-Gerlach experiment and show that the weak measurement principle works for massive particles.

The second step would be to build a 2-slit experiment and map the trajectories of massive particles. Then attempt to reconstruct the quantum potential to see if it fits the Bohm prediction.

S-G using potassium atoms - 1



S-G using potassium atoms - 2




Electrons in the 2-slit experiment

Mollenstedt and Duker. Zeitschrift fur Physik, 145 S:377-397, 1956.

Mollenstedt used an electrostatic biprism as the 2-slits used in Young's experiment.

The 10 - 20keV electron beam enters at the top at “y”.

The biprism consists of a gold covered quartz fibre, “a”, diameter of 2.5 mm and 6 mm in length, set vertically and equidistant between two grounded conductors, “b”, 4 mm apart. The fibre was held at a potential of 10 V.




The working region estimated 15 μm from the fibre.

Observation of the interference fringes for electrons


Effectively the biprism produces two virtual images of the filament emitting the electrons. The interference pattern is obtained as the superposition of the electron waves arriving in the observing plane.

By adjusting the voltage on the on the fibre the interference pattern gradually comes into sharp relief.




Neutron 2-slit experiment

Zeilinger et al. Rev.Mod.Phys., 60:1067–1073, 1988.



The neutrons are selected and focused onto a slit similar to that used by Mollenstadt.

Measurement of position of neutrons



BF_3 detectors were used to scan across horizontally and the number of neutrons were counted at regular intervals.

If a weak measurement was carried out between the slits and the BF_3 detectors then it should be possible to map out the trajectories.

Conclusion

Basil has shown the history of Bohm's interpretation and the introduction of the quantum potential and its possible use in nano-technology. He also reported the objections of Bohr, Heisenberg and Pauli to the whole approach. Their objections make people think it is in some way illegitimate.

The theory of the weak values was explained in detail and its possible use in directly observing the quantum potential.

We have explained the difference between a weak and strong measurement and how particle physics uses weak measurement without realising it.

The principle of weak measurement has been observed using photons in:

- An analogue of double Stern-Gerlach experiment.
- A Young's 2-slit experiment.

We want to further explore experimentally the weak values using massive particles such as electrons and neutrons.