

# The ILC Higgs Factory

---

**Philip Burrows**

*John Adams Institute, Oxford University*

# Outline

---

- **Introduction**
- **The Large Hadron Collider + the Higgs boson**
- **The Higgs factory**
- **The International Linear Collider (ILC)**
- **Higgs physics at ILC**
- **Project implementation and timeline**

# Large Hadron Collider (LHC)

**Largest,  
highest-energy  
particle  
collider**

**CERN,  
Geneva**



# A Higgs boson?

**ATLAS**  
EXPERIMENT





<http://atlas.ch>

Run: 203602


Event: 82614360

Date: 2012-05-18


Time: 20:28:11 CEST




# A Higgs boson?



# The 2012 discovery




# The 2012 discovery



# ATLAS status

ATLAS CONF 2014 009




$\sqrt{s} = 7 \text{ TeV} \int L dt = 4.6\text{-}4.8 \text{ fb}^{-1}$

$\sqrt{s} = 8 \text{ TeV} \int L dt = 20.3 \text{ fb}^{-1}$

Signal strength ( $\mu$ )

# It's officially a Higgs Boson!



# Finger-printing the Higgs boson

---

Determine its 'profile':

- **Mass**
- **Width**
- **Spin**
- **CP nature**
- **Coupling to fermions**
- **Coupling to gauge bosons**
- **Yukawa coupling to top quark**
- **Self coupling  $\rightarrow$  Higgs potential**

# Finger-printing the Higgs boson

---

**Is it:**

**the Higgs Boson of the Standard Model?**

**another type of Higgs boson?**

**something that looks like a Higgs boson but is actually more complicated?**

# Finger-printing the Higgs boson

---

**Is it:**

**the Higgs Boson of the Standard Model?**

**another type of Higgs boson?**

**something that looks like a Higgs boson but is actually more complicated?**

**→ Measurements of the Higgs couplings to the different species of quarks, leptons and gauge bosons are the key to answering these questions**

# Non-Standard Higgs couplings

---

**Snowmass Higgs working group:**

**Decoupling limit:**

**If all new particles (except Higgs) are at a (high) high mass scale  $M$**

**deviations from SM predictions**

**are of order  $m_H^2 / M^2$**

# Non-Standard Higgs couplings

For  $M = 1$  TeV, deviations of couplings from SM:

Model	$\kappa_V$	$\kappa_b$	$\kappa_\gamma$
Singlet Mixing	$\sim 6\%$	$\sim 6\%$	$\sim 6\%$
2HDM	$\sim 1\%$	$\sim 10\%$	$\sim 1\%$
Decoupling MSSM	$\sim -0.0013\%$	$\sim 1.6\%$	$< 1.5\%$
Composite	$\sim -3\%$	$\sim -(3 - 9)\%$	$\sim -9\%$
Top Partner	$\sim -2\%$	$\sim -2\%$	$\sim +1\%$

# Non-Standard Higgs couplings

For  $M = 1$  TeV, deviations of couplings from SM:

Model	$\kappa_V$	$\kappa_b$	$\kappa_\gamma$
Singlet Mixing	$\sim 6\%$	$\sim 6\%$	$\sim 6\%$
2HDM	$\sim 1\%$	$\sim 10\%$	$\sim 1\%$
Decoupling MSSM	$\sim -0.0013\%$	$\sim 1.6\%$	$< 1.5\%$
Composite	$\sim -3\%$	$\sim -(3 - 9)\%$	$\sim -9\%$
Top Partner	$\sim -2\%$	$\sim -2\%$	$\sim +1\%$

Deviations in the range  $1\% \rightarrow 10\%$

# Non-Standard Higgs couplings

For  $M = 1$  TeV, deviations of couplings from SM:

Model	$\kappa_V$	$\kappa_b$	$\kappa_\gamma$
Singlet Mixing	$\sim 6\%$	$\sim 6\%$	$\sim 6\%$
2HDM	$\sim 1\%$	$\sim 10\%$	$\sim 1\%$
Decoupling MSSM	$\sim -0.0013\%$	$\sim 1.6\%$	$< 1.5\%$
Composite	$\sim -3\%$	$\sim -(3 - 9)\%$	$\sim -9\%$
Top Partner	$\sim -2\%$	$\sim -2\%$	$\sim +1\%$

Deviations in the range  $1\% \rightarrow 10\%$

$\rightarrow$  measurements must be significantly more precise to resolve such deviations

# LHC projections

---

# LHC projections

---

**Currently, typically LHC projected precisions on Higgs coupling measurements assume that:**


- **Standard Model is correct**
- **No non-Standard decay modes (total width = SM)**
- **Charm and top couplings deviate from SM by same factor**

# ATLAS projections

ATLAS Simulation Preliminary

$\sqrt{s} = 14 \text{ TeV}$ :  $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$  ;  $\int \mathcal{L} dt = 3000 \text{ fb}^{-1}$


ATL PHYS PUB 2013 014



# CMS projections

$L$ ( $\text{fb}^{-1}$ )	$\kappa_\gamma$	$\kappa_W$	$\kappa_Z$	$\kappa_g$	$\kappa_b$	$\kappa_t$	$\kappa_\tau$	$\kappa_{Z\gamma}$	$\kappa_{\mu\mu}$	$\text{BR}_{\text{SM}}$
300	[5, 7]	[4, 6]	[4, 6]	[6, 8]	[10, 13]	[14, 15]	[6, 8]	[41, 41]	[23, 23]	[14, 18]
3000	[2, 5]	[2, 5]	[2, 4]	[3, 5]	[4, 7]	[7, 10]	[2, 5]	[10, 12]	[8, 8]	[7, 11]

## CMS Projection



**CMS-NOTE-2013-002**

**Yurii Maravin, LHCC Dec 2013**

# LHC projections

---

**Currently, typically LHC projected precisions on Higgs coupling measurements assume that:**

- **Standard Model is correct**
- **No non-Standard decay modes (total width = SM)**
- **Charm and top couplings deviate from SM by same factor**

**Such assumptions are not necessary for Higgs coupling measurements at e+e- Higgs Factory ...**

# 'Higgs factory'

---

- **e+e- collider:**
  - linear collider**
  - storage ring**
- **photon-photon collider:**
  - usually considered as add-on to linear collider**
- **muon collider:**
  - usually considered as a next step beyond a future neutrino factory**

# e+e- Higgs factory


e+e- annihilations:

$E > 91 + 125 = 216 \text{ GeV}$

$E \sim 250 \text{ GeV}$

$E > 91 + 250 = 341 \text{ GeV}$

$E \sim 500 \text{ GeV}$



# **e+e- colliders**

---

- **Produce annihilations of point-like particles under controlled conditions:**

# e+e- colliders

---

- **Produce annihilations of point-like particles under controlled conditions:**

**well defined centre of mass energy:  $2E$**

# e+e- colliders

---

- **Produce annihilations of point-like particles under controlled conditions:**

**well defined centre of mass energy:  $2E$**

**complete control of event kinematics:**

$$\mathbf{p} = 0, M = 2E$$

# e+e- colliders

---

- **Produce annihilations of point-like particles under controlled conditions:**


**well defined centre of mass energy:  $2E$**

**complete control of event kinematics:**

$$\mathbf{p} = 0, M = 2E$$

**polarised beam(s)**

# $e^+e^-$ annihilations



# e+e- colliders

---

- **Produce annihilations of point-like particles under controlled conditions:**

**well defined centre of mass energy:  $2E$**



**complete control of event kinematics:**

$$\mathbf{p} = 0, M = 2E$$



**polarised beam(s)**

**clean experimental environment**



# e+e- annihilations




# e+e- annihilations




# e+e- annihilations





# e+e- annihilations





$2E > 216 \text{ GeV}$



# e+e- annihilations



# e+e- annihilations



# e+e- colliders

	ILC	ILC	ILC	CLIC	CLIC	CLIC	LEP3
$\sqrt{s}$ [GeV]	250	500	1000	500	1500	3000	240
Luminosity [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	0.75	1.8	4.9	1.3	3.7	5.9	1 per IP
>0.99 $\sqrt{s}$ fraction	87%	58%	45%	54%	38%	34%	100%
polarization $e^-$	80%	80%	80%	80%	80%	80%	-
polarization $e^+$	30%	30%	20%	>50%?	>50%?	>50%?	-
beam size $\sigma_x$ [nm]	729	474	335	100	60	40	71000
beam size $\sigma_y$ [nm]	7.7	5.9	2.7	2.6	1.5	1	320
Power [MW]	128	162	300	235	364	589	200

Wyatt

# e+e- colliders

	ILC	ILC	ILC	CLIC	CLIC	CLIC	LEP3
$\sqrt{s}$ [GeV]	250	500	1000	500	1500	3000	240
Luminosity [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	0.75	1.8	4.9	1.3	3.7	5.9	1 per IP
>0.99 $\sqrt{s}$ fraction	87%	58%	45%	54%	38%	34%	100%
polarization $e^-$	80%	80%	80%	80%	80%	80%	-
polarization $e^+$	30%	30%	20%	>50%?	>50%?	>50%?	-
beam size $\sigma_x$ [nm]	729	474	335	100	60	40	71000
beam size $\sigma_y$ [nm]	7.7	5.9	2.7	2.6	1.5	1	320
Power [MW]	128	162	300	235	364	589	200

# e+e- colliders

	ILC	ILC	ILC	CLIC	CLIC	CLIC	LEP3
$\sqrt{s}$ [GeV]	250	500	1000	500	1500	3000	240
Luminosity [ $10^{34} \text{ cm}^{-1}\text{s}^{-1}$ ]	0.75	1.8	4.9	1.3	3.7	5.9	1 per IP
>0.99 $\sqrt{s}$ fraction	87%	58%	45%	54%	38%	34%	100%
polarization $e^-$	80%	80%	80%	80%	80%	80%	-
polarization $e^+$	30%	30%	20%	>50%?	>50%?	>50%?	-
beam size $\sigma_x$ [nm]	729	474	335	100	60	40	71000
beam size $\sigma_y$ [nm]	7.7	5.9	2.7	2.6	1.5	1	320
Power [MW]	128	162	300	235	364	589	200

# e+e- colliders

	ILC	ILC	ILC	CLIC	CLIC	CLIC	LEP3
$\sqrt{s}$ [GeV]	250	500	1000	500	1500	3000	240
Luminosity [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	0.75	1.8	4.9	1.3	3.7	5.9	1 per IP
>0.99 $\sqrt{s}$ fraction	87%	58%	45%	54%	38%	34%	100%
polarization $e^-$	80%	80%	80%	80%	80%	80%	-
polarization $e^+$	30%	30%	20%	>50%?	>50%?	>50%?	-
beam size $\sigma_x$ [nm]	729	474	335	100	60	40	71000
beam size $\sigma_y$ [nm]	7.7	5.9	2.7	2.6	1.5	1	320
Power [MW]	128	162	300	235	364	589	200

# e+e- colliders

	ILC	ILC	ILC	CLIC	CLIC	CLIC	LEP3
$\sqrt{s}$ [GeV]	250	500	1000	500	1500	3000	240
Luminosity [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	0.75	1.8	4.9	1.3	3.7	5.9	1 per IP
>0.99 $\sqrt{s}$ fraction	87%	58%	45%	54%	38%	34%	100%
polarization e <sup>-</sup>	80%	80%	80%	80%	80%	80%	-
polarization e <sup>+</sup>	30%	30%	20%	>50%?	>50%?	>50%?	-
beam size $\sigma_x$ [nm]	729	474	335	100	60	40	71000
beam size $\sigma_y$ [nm]	7.7	5.9	2.7	2.6	1.5	1	320
Power [MW]	128	162	300	235	364	589	200

**$L \sim 10^{34}$  (250 GeV)  $\rightarrow$  20,000 H / year**

# e+e- colliders

	ILC	ILC	ILC	CLIC	CLIC	CLIC	LEP3
$\sqrt{s}$ [GeV]	250	500	1000	500	1500	3000	240
Luminosity [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	0.75	1.8	4.9	1.3	3.7	5.9	1 per IP
>0.99 $\sqrt{s}$ fraction	87%	58%	45%	54%	38%	34%	100%
polarization e <sup>-</sup>	80%	80%	80%	80%	80%	80%	-
polarization e <sup>+</sup>	30%	30%	20%	>50%?	>50%?	>50%?	-
beam size $\sigma_x$ [nm]	729	474	335	100	60	40	71000
beam size $\sigma_y$ [nm]	7.7	5.9	2.7	2.6	1.5	1	320
Power [MW]	128	162	300	235	364	589	200

# e+e- colliders

	ILC	ILC	ILC	CLIC	CLIC	CLIC	LEP3
$\sqrt{s}$ [GeV]	250	500	1000	500	1500	3000	240
Luminosity [ $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ ]	0.75	1.8	4.9	1.3	3.7	5.9	1 per IP
>0.99 $\sqrt{s}$ fraction	87%	58%	45%	54%	38%	34%	100%
polarization e <sup>-</sup>	80%	80%	80%	80%	80%	80%	-
polarization e <sup>+</sup>	30%	30%	20%	>50%?	>50%?	>50%?	-
beam size $\sigma_x$ [nm]	729	474	335	100	60	40	71000
beam size $\sigma_y$ [nm]	7.7	5.9	2.7	2.6	1.5	1	320
Power [MW]	128	162	300	235	364	589	200

# European particle physics strategy 2013

---

**There is a strong scientific case for an electron-positron collider, complementary to the LHC, that can study the properties of the Higgs boson and other particles with unprecedented precision and whose energy can be upgraded.**

# European particle physics strategy 2013

---

**There is a strong scientific case for an electron-positron collider, complementary to the LHC, that can study the properties of the Higgs boson and other particles with unprecedented precision and whose energy can be upgraded.**

**The Technical Design Report of the International Linear Collider (ILC) has been completed, with large European participation. The initiative from the Japanese particle physics community to host the ILC in Japan is most welcome, and European groups are eager to participate.**

# European particle physics strategy 2013

---

**There is a strong scientific case for an electron-positron collider, complementary to the LHC, that can study the properties of the Higgs boson and other particles with unprecedented precision and whose energy can be upgraded.**

**The Technical Design Report of the International Linear Collider (ILC) has been completed, with large European participation. The initiative from the Japanese particle physics community to host the ILC in Japan is most welcome, and European groups are eager to participate.**

***Europe looks forward to a proposal from Japan to discuss a possible participation.***

# Snowmass executive summary 2013

---

Compelling science motivates continuing this program with experiments at lepton colliders. Experiments at such colliders can reach sub-percent precision in Higgs boson properties in a unique, model-independent way, enabling discovery of percent-level deviations from the Standard Model predicted in many theories.

# Snowmass executive summary 2013

---



Compelling science motivates continuing this program with experiments at lepton colliders. Experiments at such colliders can reach sub-percent precision in Higgs boson properties in a unique, model-independent way, enabling discovery of percent-level deviations from the Standard Model predicted in many theories. **They can improve the precision of our knowledge of the W, Z, and top quark well enough to allow the discovery of predicted new-physics effects. They search for new particles in a manner complementing new particle searches at the LHC.**

# Snowmass executive summary 2013

---

Compelling science motivates continuing this program with experiments at lepton colliders. Experiments at such colliders can reach sub-percent precision in Higgs boson properties in a unique, model-independent way, enabling discovery of percent-level deviations from the Standard Model predicted in many theories. They can improve the precision of our knowledge of the W, Z, and top quark well enough to allow the discovery of predicted new-physics effects. They search for new particles in a manner complementing new particle searches at the LHC. **A global effort has completed the technical design of the International Linear Collider (ILC) accelerator and detectors that will provide these capabilities in the latter part of the next decade. The Japanese particle physics community has declared this facility as its first priority for new initiatives.**

# $e^+e^-$ Higgs Factory



# ILC Higgs Factory Roadmap

---

## 250 GeV:

Mass, Spin, CP nature

Absolute meas. of HZZ

BRs Higgs  $\rightarrow$  qq, ll, VV

## 350 GeV:

Top threshold: mass, width, anomalous couplings ...

(more stats on Higgs BRs)

## 500 GeV:


HWW coupling  $\rightarrow$  total width  $\rightarrow$  absolute couplings

Higgs self coupling

Top Yukawa coupling

$\rightarrow$  1000 GeV: as motivated by physics


# Higgs mass measurement



$$M_X^2 = (p_{CM} - (p_{\mu^+} + p_{\mu^-}))^2$$

**Recoil mass:**  
 - independent of  
 Higgs decay

**Discovery mode**  
 for 'H' decay to  
 weakly-interacting  
 particles



250 fb<sup>-1</sup> @ 250 GeV

$$\Delta\sigma_H / \sigma_H = 2.5\%$$


$$\Delta m_H = 30 \text{ MeV}$$

(Fujii)


# Higgs spin determination

Rise of  
cross-section  
near threshold

(TESLA TDR)



# Higgs branching ratios determination (1)



ILD DBD Study (Ono)


$250 \text{ fb}^{-1} @ 250 \text{ GeV}$   
 $m_H = 120 \text{ GeV}$

	@ 250 GeV
process	ZH
luminosity · fb	250
cross section	2.5%
	$\sigma \cdot \text{Br}$
H → bb	1.0%
H → cc	6.9%
H → gg	8.5%
H → WW*	8.2%
H → ττ	4-6%
H → ZZ*	28%
H → γγ	23-30%

(ILC TDR)

# Higgs branching ratios determination (2)

measurements (independent)	precision
$X_1 = \sigma_{ZH} \cdot \text{Br}(H \rightarrow b\bar{b}) @ 250 \text{ GeV}$	1.0%
$X_2 = \sigma_{ZH} \cdot \text{Br}(H \rightarrow c\bar{c}) @ 250 \text{ GeV}$	6.9%
$X_3 = \sigma_{ZH} \cdot \text{Br}(H \rightarrow gg) @ 250 \text{ GeV}$	8.5%
$X_4 = \sigma_{ZH} \cdot \text{Br}(H \rightarrow WW^*) @ 250 \text{ GeV}$	8.2%
$X_5 = \sigma_{ZH} \cdot \text{Br}(H \rightarrow b\bar{b}) @ 500 \text{ GeV}$	1.6%
$X_6 = \sigma_{ZH} \cdot \text{Br}(H \rightarrow c\bar{c}) @ 500 \text{ GeV}$	11%
$X_7 = \sigma_{ZH} \cdot \text{Br}(H \rightarrow gg) @ 500 \text{ GeV}$	13%
$X_8 = \sigma_{\nu\bar{\nu}H} \cdot \text{Br}(H \rightarrow b\bar{b}) @ 500 \text{ GeV}$	0.60%
$X_9 = \sigma_{\nu\bar{\nu}H} \cdot \text{Br}(H \rightarrow c\bar{c}) @ 500 \text{ GeV}$	4.0%
$X_{10} = \sigma_{\nu\bar{\nu}H} \cdot \text{Br}(H \rightarrow gg) @ 500 \text{ GeV}$	4.9%
$X_{11} = \sigma_{\nu\bar{\nu}H} \cdot \text{Br}(H \rightarrow WW^*) @ 500 \text{ GeV}$	3.0%
$X_{12} = \sigma_{ZH}$	2.5%



(Fujii / ILC TDR)

# Total Width and Coupling Extraction

One of the major advantages of the LC

To extract couplings from BRs, we need the total width:

$$g_{HAA}^2 \propto \Gamma(H \rightarrow AA) = \Gamma_H \cdot BR(H \rightarrow AA)$$

To determine the total width, we need at least one partial width and corresponding BR:

$$\Gamma_H = \Gamma(H \rightarrow AA) / BR(H \rightarrow AA)$$

In principle, we can use the  $A=Z$ , or  $W$  for which we can measure both the BRs and the couplings:

A Feynman diagram showing the production and decay of a Higgs boson (H) via a Z boson. An electron-positron pair ( $e^+e^-$ ) annihilates into a Z boson. The Z boson then decays into a Higgs boson (H) and another Z boson. The Higgs boson subsequently decays into a Z boson and an anti-Z boson ( $ZZ^*$ ). A red dashed line indicates the Higgs boson, and a black arrow points to the vertex where the Higgs is produced. The diagram is labeled with  $\Gamma(H \rightarrow ZZ^*)$  and  $BR(H \rightarrow ZZ^*)$ .



BR=O(1%): precision limited by low stat.  
for  $H \rightarrow ZZ^*$  events

A Feynman diagram showing the production and decay of a Higgs boson (H) via a W boson. An electron-positron pair ( $e^+e^-$ ) annihilates into a neutrino-antineutrino pair ( $\bar{\nu}\nu$ ) and a W boson. The W boson then decays into a Higgs boson (H) and another W boson. The Higgs boson subsequently decays into a W boson and an anti-W boson ( $WW^*$ ). A red dashed line indicates the Higgs boson, and a black arrow points to the vertex where the Higgs is produced. The diagram is labeled with  $\Gamma(H \rightarrow WW^*)$  and  $BR(H \rightarrow WW^*)$ .

More advantageous but not easy at low E  
250 fb<sup>-1</sup>@250 GeV  
 $\Delta\Gamma_H/\Gamma_H \simeq 10\%$



Jenny List's talk

# Higgs self-coupling determination



$\sqrt{s}$ (GeV)	500	500	500+1000	500+1000
L ( $\text{fb}^{-1}$ )	500	1600	500+1000	1600+2500
$\Delta\lambda/\lambda$	83%	46%	21%	13%

# Higgs top-coupling determination



$1 \text{ ab}^{-1} @ 500 \text{ GeV}$

$$\Delta g_Y(t) / g_Y(t) = 10 \%$$

(Price, Roloff)

# ILC roadmap

---

<b>Baseline:</b>	<b>250 fb<sup>-1</sup></b>	<b>@ 250 GeV</b>	<b>3 years</b>
	<b>500 fb<sup>-1</sup></b>	<b>@ 500 GeV</b>	<b>3 years</b>
	<b>1000 fb<sup>-1</sup></b>	<b>@ 1000 GeV</b>	<b>3 years</b>

# ILC roadmap

---

<b>Baseline:</b>	<b>250 fb<sup>-1</sup></b>	<b>@ 250 GeV</b>	<b>3 years</b>
	<b>500 fb<sup>-1</sup></b>	<b>@ 500 GeV</b>	<b>3 years</b>
	<b>1000 fb<sup>-1</sup></b>	<b>@ 1000 GeV</b>	<b>3 years</b>


**Followed by luminosity upgrade:**

<b>'HL-ILC':</b>	<b>+900 fb<sup>-1</sup></b>	<b>@ 250 GeV</b>	<b>+3 years</b>
	<b>+1100 fb<sup>-1</sup></b>	<b>@ 500 GeV</b>	<b>+3 years</b>
	<b>+1500 fb<sup>-1</sup></b>	<b>@ 1000 GeV</b>	<b>+3 years</b>

# ILC baseline precisions

$\sqrt{s}$ and $\mathcal{L}$ ( $P_{e^-}, P_{e^+}$ )	250 fb <sup>-1</sup> at 250 GeV (-0.8, +0.3)		500 fb <sup>-1</sup> at 500 GeV (-0.8, +0.3)				1 ab <sup>-1</sup> at 1 TeV (-0.8, +0.2)		
	$Zh$	$\nu\bar{\nu}h$	$Zh$	$\nu\bar{\nu}h$	$t\bar{t}h$	$Zhh$	$\nu\bar{\nu}h$	$t\bar{t}h$	$\nu\bar{\nu}hh$
$\Delta\sigma/\sigma$	2.6%	-	3.0	-		42.7%			26.3%
BR(invis.)	< 0.9 %	-	-	-	-				
mode	$\Delta(\sigma \cdot BR)/(\sigma \cdot BR)$								
$h \rightarrow bb$	1.2%	<b>10.5%</b>	1.8%	0.7%	28%		0.5%	<b>6.0%</b>	
$h \rightarrow c\bar{c}$	8.3%	-	13%	6.2%			3.1%		
$h \rightarrow gg$	7.0%	-	11%	4.1%			2.3%		
$h \rightarrow WW^*$	6.4%	-	9.2%	2.4%			1.6%		
$h \rightarrow \tau^+\tau^-$	4.2%	-	5.4%	9.0%			3.1%		
$h \rightarrow ZZ^*$	19%	-	25%	8.2%			4.1%		
$h \rightarrow \gamma\gamma$	34%	-	34%	23%			8.5%		
$h \rightarrow \mu^+\mu^-$	100%	-	-	-			31%		

# Higgs coupling map



(Fujii)

# ILC baseline + HL-ILC precisions

$\sqrt{s}$ and $\mathcal{L}$ ( $P_{e^-}, P_{e^+}$ )	1150 fb <sup>-1</sup> at 250 GeV (-0.8, +0.3)		1600 fb <sup>-1</sup> at 500 GeV (-0.8, +0.3)				2.5 ab <sup>-1</sup> at 1 TeV (-0.8, +0.2)		
	$Zh$	$\nu\bar{\nu}h$	$Zh$	$\nu\bar{\nu}h$	$t\bar{t}h$	$Zhh$	$\nu\bar{\nu}h$	$t\bar{t}h$	$\nu\bar{\nu}hh$
$\Delta\sigma/\sigma$	1.2%	-	1.7	-		23.7%			16.7%
BR(invis.)	< 0.4 %	-	-	-			-		

mode	$\Delta(\sigma \cdot BR)/(\sigma \cdot BR)$								
$h \rightarrow b\bar{b}$	0.6%	4.9%	1.0%	0.4%	16%		0.3%	3.8%	
$h \rightarrow c\bar{c}$	3.9%	-	7.2%	3.5%			2.0%		
$h \rightarrow gg$	3.3%	-	6.0%	2.3%			1.4%		
$h \rightarrow WW^*$	3.0%	-	5.1%	1.3%			1.0%		
$h \rightarrow \tau^+\tau^-$	2.0%	-	3.0%	5.0%			2.0%		
$h \rightarrow ZZ^*$	8.8%	-	14%	4.6%			2.6%		
$h \rightarrow \gamma\gamma$	16%	-	19%	13%			5.4%		
$h \rightarrow \mu^+\mu^-$	46.6%	-	-	-			20%		

# Model-independent couplings extraction

33 input measurements

11-parameter fit

$$\chi^2 = \sum_{i=1}^{i=33} \left( \frac{Y_i - Y'_i}{\Delta Y_i} \right)^2,$$


$$Y'_i = F_i \cdot \frac{g_{HZZ}^2 g_{Hb\bar{b}}^2}{\Gamma_0}, \text{ or } Y'_i = F_i \cdot \frac{g_{HWW}^2 g_{Hb\bar{b}}^2}{\Gamma_0}, \text{ or } Y'_i = F_i \cdot \frac{g_{Htt}^2 g_{Hb\bar{b}}^2}{\Gamma_0}$$

$$F_i = S_i G_i \quad \text{where } S_i = \left( \frac{\sigma_{ZH}}{g_Z^2} \right), \left( \frac{\sigma_{\nu\bar{\nu}H}}{g_W^2} \right), \text{ or } \left( \frac{\sigma_{t\bar{t}H}}{g_t^2} \right), \text{ and } G_i = \left( \frac{\Gamma_i}{g_i^2} \right).$$

# Model-independent couplings

	ILC(250)	ILC(500)	ILC(1000)	ILC(LumUp)
$\sqrt{s}$ (GeV)	250	250+500	250+500+1000	250+500+1000
L ( $\text{fb}^{-1}$ )	250	250+500	250+500+1000	1150+1600+2500
$\gamma\gamma$	18 %	8.4 %	4.0 %	2.4 %
$gg$	6.4 %	2.3 %	1.6 %	0.9 %
$WW$	4.8 %	1.1 %	1.1 %	0.6 %
$ZZ$	1.3 %	1.0 %	1.0 %	0.5 %
$t\bar{t}$	–	14 %	3.1 %	1.9 %
$b\bar{b}$	5.3 %	1.6 %	1.3 %	0.7 %
$\tau^+\tau^-$	5.7 %	2.3 %	1.6 %	0.9 %
$c\bar{c}$	6.8 %	2.8 %	1.8 %	1.0 %
$\mu^+\mu^-$	91%	91%	16 %	10 %
$\Gamma_T(h)$	12 %	4.9 %	4.5 %	2.3 %
$hhh$	–	83 %	21 %	13 %
BR(invis.)	< 0.9 %	< 0.9 %	< 0.9 %	< 0.4 %

# Model-independent couplings



# Comparison with LHC

---

**LHC does not project making model-independent Higgs coupling measurements**

**LHC projections assume the Standard Model and estimate precision relative to SM couplings, also assuming charm follows top**

# Model-dependent couplings extraction

7 Parameter HXSWG Benchmark \*

LHC

Mode	LHC	
	300 fb <sup>-1</sup>	3000 fb <sup>-1</sup>
$\gamma\gamma$	(5 – 7)%	(2 – 5)%
$gg$	(6 – 8)%	(3 – 5)%
$WW$	(4 – 5)%	(2 – 3)%
$ZZ$	(4 – 5)%	(2 – 3)%
$t\bar{t}$	(14 – 15)%	(7 – 10)%
$b\bar{b}$	(10 – 13)%	(4 – 7)%
$\tau^+\tau^-$	(6 – 8)%	(2 – 5)%

\* Assume  $\kappa_c = \kappa_t$  &  $\Gamma_{tot} = \sum_{\text{SM decays } i} \Gamma_i^{SM} \kappa_i^2$

# Comparison with LHC

---

**LHC does not project making model-independent Higgs coupling measurements**

**LHC projections assume the Standard Model and estimate precision relative to SM couplings, also assuming charm follows top**

**For purpose of comparison, can follow same model-dependent procedure for ILC ...**

# Model-dependent couplings extraction

7 Parameter HXSWG Benchmark \*

Mode	LHC		ILC(1000)	ILC(LumUp)	$\sqrt{s}$ (GeV) L ( $\text{fb}^{-1}$ )
	300 $\text{fb}^{-1}$	3000 $\text{fb}^{-1}$	250+500+1000	250+500+1000	
			250+500+1000	1150+1600+2500	
$\gamma\gamma$	(5 – 7)%	(2 – 5)%	3.8 %	2.3 %	
$gg$	(6 – 8)%	(3 – 5)%	1.1 %	0.7 %	
$WW$	(4 – 5)%	(2 – 3)%	0.3 %	0.2 %	
$ZZ$	(4 – 5)%	(2 – 3)%	0.5 %	0.3 %	
$t\bar{t}$	(14 – 15)%	(7 – 10)%	1.3 %	0.9 %	
$b\bar{b}$	(10 – 13)%	(4 – 7)%	0.6 %	0.4 %	
$\tau^+\tau^-$	(6 – 8)%	(2 – 5)%	1.3 %	0.7 %	

\* Assume  $\kappa_c = \kappa_t$  &  $\Gamma_{tot} = \sum_{\text{SM decays } i} \Gamma_i^{SM} \kappa_i^2$

# Model-dependent couplings extraction

7 Parameter HXSWG Benchmark \*

Mode	LHC		ILC(1000)	ILC(LumUp)	$\sqrt{s}$ (GeV) L ( $\text{fb}^{-1}$ )
	300 $\text{fb}^{-1}$	3000 $\text{fb}^{-1}$	250+500+1000 250+500+1000	250+500+1000 1150+1600+2500	
$\gamma\gamma$	(5 – 7)%	(2 – 5)%	3.8 %	2.3 %	
$gg$	(6 – 8)%	(3 – 5)%	1.1 %	0.7 %	
$WW$	(4 – 5)%	(2 – 3)%	0.3 %	0.2 %	
$ZZ$	(4 – 5)%	(2 – 3)%	0.5 %	0.3 %	
$t\bar{t}$	(14 – 15)%	(7 – 10)%	1.3 %	0.9 %	
$b\bar{b}$	(10 – 13)%	(4 – 7)%	0.6 %	0.4 %	
$\tau^+\tau^-$	(6 – 8)%	(2 – 5)%	1.3 %	0.7 %	

**~10 x LHC sensitivity**

\* Assume  $\kappa_c = \kappa_t$  &  $\Gamma_{tot} = \sum_{\text{SM decays } i} \Gamma_i^{SM} \kappa_i^2$

# Non-Standard Higgs couplings

For  $M = 1$  TeV, deviations of couplings from SM:


Model	$\kappa_V$	$\kappa_b$	$\kappa_\gamma$
Singlet Mixing	$\sim 6\%$	$\sim 6\%$	$\sim 6\%$
2HDM	$\sim 1\%$	$\sim 10\%$	$\sim 1\%$
Decoupling MSSM	$\sim -0.0013\%$	$\sim 1.6\%$	$< 1.5\%$
Composite	$\sim -3\%$	$\sim -(3 - 9)\%$	$\sim -9\%$
Top Partner	$\sim -2\%$	$\sim -2\%$	$\sim +1\%$

Deviations in the range  $1\% \rightarrow 10\%$


**$\rightarrow$  measurements must be significantly more precise to resolve such deviations**

# Specific beyond-SM examples

## Composite Higgs (MCHM5)



Zivkovic et al



Simulated ILC measurements

# The accelerator

---

# Large Electron Positron collider (RIP)



**0.1 TeV  
beams**

# Large Electron Positron collider (RIP)



**0.1 TeV  
beams**

**Synch**

**rad →**

**18 MW**

# Super Large Electron Positron collider?



**0.2 TeV**  
**beams?**

# Super Large Electron Positron collider?



**0.2 TeV**  
**beams?**

**Synch**  
**rad →**  
**300 MW**

# Super Large Electron Positron collider?



**0.2 TeV**  
**beams?**

**Synch**

**rad →**


**300 MW**

# Linear Colliders for electrons + positrons

**Stanford  
Linear  
Accelerator  
Center  
(California)**





# Designing a Linear Collider



# International Linear Collider (ILC)

- RTML
- ML
- SOURCES & BDS
- DR
- DETECTOR AREA

c. 250 GeV / beam



31 km




# Beam parameters

---

## ILC (500)

<b>Electrons/bunch</b>	<b>0.75</b>	<b>10**10</b>
<b>Bunches/train</b>	<b>2820</b>	
<b>Train repetition rate</b>	<b>5</b>	<b>Hz</b>
<b>Bunch separation</b>	<b>308</b>	<b>ns</b>
<b>Train length</b>	<b>868</b>	<b>us</b>
<b>Horizontal IP beam size</b>	<b>655</b>	<b>nm</b>
<b>Vertical IP beam size</b>	<b>6</b>	<b>nm</b>
<b>Longitudinal IP beam size</b>	<b>300</b>	<b>um</b>
<b>Luminosity</b>	<b>2</b>	<b>10**34</b>

# ILC Detectors



# ILC project status

---


- **2005-12 ILC run by Global Design Effort (Barish)**
- **C. 500 accelerator scientists worldwide involved**
- **A Reference Design Report (RDR) was completed in 2007**  
**including a first cost estimate**
- **2008-12 engineering design phase**  
**major focus on risk minimisation + cost reduction**
- **Technical Design document released end 2012**  
**revised cost estimate + project implementation plan**

# Technical Volumes

2007

2011

2013



Reference Design Report




ILC Technical Progress Report ("interim report")

AD&I



~250 pages  
Deliverable 2



~300 pages  
Deliverables 1,3 and 4

Technical Design Report

# ILC project status

---

- **2005-12 ILC run by Global Design Effort (Barish)**
- **C. 500 accelerator scientists worldwide involved**
- **A Reference Design Report (RDR) was completed in 2007**  
**including a first cost estimate**
- **2008-12 engineering design phase**  
**major focus on risk minimisation + cost reduction**
- **Technical Design document released end 2012**  
**revised cost estimate + project implementation plan**
- **Lyn Evans assumed project leadership 2013**  
**Japan preparing implementation of ILC at Kitakami**

# ILC Plan in Japan


- ▶ Japanese HEP community proposes to host ILC based on the “staging scenario” to the Japanese Government.
  - ILC starts as a 250GeV Higgs factory, and will evolve to a 500GeV machine.
  - Technical extendability to 1TeV is to be preserved.

# ILC Plan in Japan

- ▶ Japanese HEP community proposes to host ILC based on the "staging scenario" to the Japanese

Government

## - Japanese Mountainous Sites -



# LDP (Liberal Democratic Party) Victory in the lower-house election in Oct, 2012

Our new prime minister  
Shinzo Abe



LDP took power in Dec 2012

The ILC appears twice explicitly in  
the policy document:

- Science and technology policies
- Creation of top-class research centers

LDP policy document  
for the election

**Yamamoto, HEPAP, 11/3/13**

# ILC in Japan?



**meeting of Lyn Evans and Prime Minister Abe, March 27, 2013**

# Possible Timeline

- July 2013
  - Non-political evaluation of 2 Japanese candidate sites complete, followed by down-selecting to one
- End 2013
  - Japanese government announces its intent to bid
- 2013~2015
  - Inter-governmental negotiations
  - Completion of R&Ds, preparation for the ILC lab.
- ~2015
  - Inputs from LHC@14TeV, decision to proceed
- 2015~16
  - Construction begins (incl. bidding)
- 2026~27
  - Commissioning

# PPAP recommendation


## The UK Particle Physics Roadmap

Particle Physics Advisory Panel:

*P. N. Burrows, C. Da Via, E. W. N. Glover, P.R. Newman, J. Rademacker,  
C. Shepherd-Themistocleous, W.J. Spence, M. A. Thomson and M. Wing*

7/11/12

**‘It is essential that the UK engages with the Higgs Factory initiative and positions itself to play a leading role should the facility go ahead.’**



# Extra material follows

---

# Model-independent couplings

Facility	ILC			ILC(LumiUp)	TLEP (4 IP)		CLIC		
$\sqrt{s}$ (GeV)	250	500	1000	250/500/1000	240	350	350	1400	3000
$\int \mathcal{L} dt$ (fb $^{-1}$ )	250	+500	+1000	1150+1600+2500 $^\ddagger$	10000	+2600	500	+1500	+2000
$P(e^-, e^+)$	(-0.8, +0.3)	(-0.8, +0.3)	(-0.8, +0.2)	(same)	(0, 0)	(0, 0)	(0, 0)	(-0.8, 0)	(-0.8, 0)
$\Gamma_H$	12%	5.0%	4.6%	2.5%	1.9%	1.0%	9.2%	8.5%	8.4%
$\kappa_\gamma$	18%	8.4%	4.0%	2.4%	1.7%	1.5%	–	5.9%	<5.9%
$\kappa_g$	6.4%	2.3%	1.6%	0.9%	1.1%	0.8%	4.1%	2.3%	2.2%
$\kappa_W$	4.9%	1.2%	1.2%	0.6%	0.85%	0.19%	2.6%	2.1%	2.1%
$\kappa_Z$	1.3%	1.0%	1.0%	0.5%	0.16%	0.15%	2.1%	2.1%	2.1%
$\kappa_\mu$	91%	91%	16%	10%	6.4%	6.2%	–	11%	5.6%
$\kappa_\tau$	5.8%	2.4%	1.8%	1.0%	0.94%	0.54%	4.0%	2.5%	<2.5%
$\kappa_c$	6.8%	2.8%	1.8%	1.1%	1.0%	0.71%	3.8%	2.4%	2.2%
$\kappa_b$	5.3%	1.7%	1.3%	0.8%	0.88%	0.42%	2.8%	2.2%	2.1%
$\kappa_t$	–	14%	3.2%	2.0%	–	13%	–	4.5%	<4.5%
$BR_{inv}$	0.9%	< 0.9%	< 0.9%	0.4%	0.19%	< 0.19%			

Oct 28, 2014 21

# Key challenges


- **Energy:**  
sustain high gradients  
> 30 MeV/m
- **Luminosity:**



# Niobium Accelerating Cavities



**TM010 mode**



# Niobium Accelerating Cavities




**c. 20,000 needed**


# Niobium Accelerating Cavities



**c. 20,000 needed**




# Niobium Accelerating Cavities



# European X-FEL at DESY

← 3.4km →



# Key challenges

- **Energy:**  
sustain high gradients  
> 30 MeV/m
- **Luminosity:**  
goal is  
>  $10^{34}$  / cm<sup>2</sup>/ s



# ILC Cost Estimate (February 2007)

---

- **shared value = 4.87 Billion ILC Value Units**
- **site-dependent value = 1.78 Billion ILC Value Units**
- **total value = 6.65 Billion ILC Value Units**  
*(shared + site-dependent)*
  
- **labour = 22 million person-hours = 13,000 person-years**  
*(assuming 1700 person-hours per person-year)*

**1 ILC Value Unit = 1 US Dollar (2007) = 0.83 Euros = 117 Yen**

# ILC value breakdown

